



DALBAVANCIN VARIABLE DOSE

Patient details Name Address NHS number DOB			No known a	allergies ature, details and date of	Indication for treatment: Date antibiotic to start in community: Planned treatment length in community:	Consultant/ G	Consultant/ GP Practice:					
eGFR:	C	reatinine:		date:	Weight (kg): date							
Medication	Dose	Frequency	/ Route	Instructions for preparation and use								
DALBAVANCIN		STAT	IV	Reconstitute 500 mg Dalbavancin vial(s) by adding 25mL water for injections slowly to each vial, directing the flow down the vial wall. To avoid foaming, gently swirl and invert the vial until contents dissolved (may take up to 5 minutes). Do not shake. TransfermL of the reconstituted dose to amlL glucose 5% bag.								
Water for Injections (For reconstitution)	mL	STAT	IV	Administration: Give via IV infusion 30 minutes using a pump. ^{1&2}								
Glucose 5% (Diluent)	mL	STAT	IV									
Sodium Chloride 0.9% (For Infusion Set Flush)	As SOP ³	OD	IV	Agilia Volumetric Pump	b : Administer 25 mL at the same rate as the infusion above .							
Sodium Chloride 0.9%	Flush the c	Flush the cannula with 5mL of sodium chloride 0.9% or the PICC line with 10mL sodium chloride 0.9% before and after each administration via IV route.										
, .				•	e from: https://www.medicines.org.uk/emc/product/2270/smpc. 2 .Medusa. Dalbavancin. I y.asp. 3. SOP for Intravenous Infusion Set Flushing – available through NBT/UHBW link.	travenous infusion. In	jectable					

Date & time:											
Given by:											

Signed:	Name:	Professional	Bleep/	Date:	
	(Print Name)	registration number:	Telephone:		