

## Adult Community IV Antibiotic Treatment: Authorisation to Administer and Administration Record: Flucloxacillin 8g elastomeric pump

<b>Patient details</b> Name Address  NHS number DOB	<b>Allergies &amp; Intolerances:</b>  No known allergies <input type="checkbox"/> Document nature, details and date of each reaction	<b>Indication for treatment:</b>  Date antibiotic to start in community:  Planned treatment length in community or end date:			
<b>eGFR:</b>	<b>Creatinine:</b>	<b>Date:</b>	<b>Weight (kg):</b>	<b>Date:</b>	
Medication	Dose	Frequency	Route	Instructions for preparation and use	Pharmacy check
Flucloxacillin (unlicensed)	8g (in 230mL Baxter LV10 elastomeric device)	Continuous infusion over 24 hours	IV	<b>Administration:</b> Connect the elastomeric device as per guidelines. Once connected the elastomeric device is designed to administer the dose by continuous infusion over 24 hours	
Sodium Chloride 0.9%	Flush PICC line with 10mL sodium chloride 0.9% before each dose and at the end of the treatment course				

Date & time:	/	/	/	/	/	/	/	/	/	/	/	/	/	/	/	/	/	/	/	
Given by:																				

Prescriber must be F2 or above, or a suitable non-medical prescriber.									
Signed:		Name: (Print Name)		Professional registration number:		Bleep/ Telephone:		Date:	

