

Adult Community IV Antibiotic Treatment: Authorisation to Administer and Administration Record
MEROPENEM 500mg TDS

Patient details Name Address NHS number DOB	Allergies & Intolerances: No known allergies <input type="checkbox"/> Document nature, details and date of each reaction	Indication for treatment: Date antibiotic to start in community: Planned treatment length in community or end date:			
eGFR:	Creatinine:	Date:	Weight (kg):	Date:	
Medication	Dose	Frequency	Route	Instructions for preparation and use	Pharmacy check
Meropenem	500mg	TDS	IV	Reconstitute a 500mg meropenem vial with 10mL of water for injections. Shake well until the solution is clear (this may take some time). Reconstituted solutions should be clear, colourless to yellow and particulate free. Administration: Give the 500mg slowly over 5 minutes ¹⁻²	
Water for injections	10mL	TDS	IV		
Sodium Chloride 0.9%	Flush the cannula with 5ml of sodium chloride 0.9% or the PICC line with 10mL sodium chloride 0.9% before and after each administration via IV route.				

1. SPC. Meropenem 500mg powder for solution for injection/ infusion. Last updated March 2021. Available from <https://www.medicines.org.uk/emc/medicine/34860/SPC#>
2. Meropenem. Intravenous. In: Injectable Medicines Guide. Last updated August 2018. Available from: <https://medusa.wales.nhs.uk/IVGuideDisplay.asp>

Date & time:																				
Given by:																				

Prescriber must be F2 or above, or a suitable non-medical prescriber.

Signed:		Name: (Print Name)		Professional registration number:		Bleep/ Telephone:		Date:	
----------------	--	------------------------------	--	--	--	--------------------------	--	--------------	--