

## Adult Community IV Antibiotic Treatment: Authorisation to Administer and Administration Record

### TEICOPLANIN VARIABLE DOSING – 800MG OR ABOVE

<b>Patient details</b> Name Address  NHS number DOB	<b>Allergies &amp; Intolerances:</b>  <b>No known allergies</b> <input type="checkbox"/> Document nature, details and date of each reaction	<b>Indication for treatment:</b>  <b>Date antibiotic to start in community:</b>  <b>Planned treatment length in community or end date:</b>
--	--	--

<b>eGFR:</b>	<b>Creatinine:</b>	<b>Date:</b>	<b>Weight (kg):</b>	<b>Date:</b>
--------------	--------------------	--------------	---------------------	--------------

Medication	Dose	Frequency	Route	Instructions for preparation and use	Pharmacy check
Teicoplanin	-----	OD	IV	Reconstitute ___ 400mg Teicoplanin vial(s) and ___ 200mg Teicoplanin vial(s) with the water for injections supplied with each of the teicoplanin vials. Avoid foam formation, do not shake vial but gently roll. If foam develops allow to stand for 15 minutes. The resultant 3mL solution contains the total stated dose of teicoplanin in the vial. Only clear, yellowish solutions should be used. A 23G needle must be used to reconstitute the vial and when withdrawing the required dose from the vial. Withdrawal ___ mL of the reconstituted solution and transfer into a 100mL sodium chloride 0.9% bag. <b>Administration:</b> Give as an infusion over 30 minutes via infusion pump.	
Water for injections (Supplied with teicoplanin vial)	___ x 3.14mL	OD	IV		
Sodium Chloride 0.9%	100mL	OD	IV		
Sodium Chloride 0.9% <i>(For Infusion Set Flush)</i>	As SOP <sup>3</sup>	OD	IV	1) <b>Agilia Volumetric Pump:</b> Administer 25 mL at the <b>same rate as the infusion above.</b> 2) <b>Gravity Infusion:</b> The flush volume is the priming volume of infusion set. Withdraw excess sodium chloride 0.9% from the bag to leave flush volume and administer at the <b>same rate as the infusion above.</b>	
Sodium Chloride 0.9%	Flush the cannula with 5ml of sodium chloride 0.9% or the PICC line with -10mL sodium chloride 0.9% before and after each administration via IV route.				

1. Medusa Available form Injectable Medicines Guide - Display - Teicoplanin - Intravenous - Version 14 - IVGuideDisplayMain.asp (wales.nhs.uk) Last updated 04/10/23. 2 SPC Available from Targocid 400mg powder for solution for injection/infusion or oral solution - Summary of Product Characteristics (SmPC) - (emc) (medicines.org.uk) Last updated 21/10/2022 3. SOP for Intravenous Infusion Set Flushing – available through NBT LINK/UHBW

<b>Date &amp; time:</b>																				
<b>Given by:</b>																				

Prescriber must be F2 or above, or a suitable non-medical prescriber.

<b>Signed:</b>		<b>Name:</b> (Print Name)		<b>Professional registration number:</b>		<b>Bleep/ Telephone:</b>		<b>Date:</b>	
----------------	--	------------------------------	--	--	--	--------------------------	--	--------------	--