

Adult Community IV Antibiotic Treatment: Authorisation to Administer and Administration Record

TEICOPLANIN 600mg OD POST LOADING

Patient details Name Address NHS number DOB		Allergies & Intolerances: No known allergies <input type="checkbox"/> Document nature, details and date of each reaction		Indication for treatment: Date antibiotic to start in community: Planned treatment length in community or end date:	
eGFR:		Creatinine:		Date:	
Weight (kg):		Date:			
Medication	Dose	Frequency	Route	Instructions for preparation and use	Pharmacy check
Teicoplanin	400mg	OD	IV	Reconstitute 1x 400mg and 1x200mg vial (or alternatively 3 x 200mg vials) with the water for injections supplied with each teicoplanin vial. Avoid foam formation, do not shake vial but gently roll. If foam develops allow to stand for 15 minutes. The resultant 3mL solution contains the total stated dose of teicoplanin in the vial. Only clear, yellowish solutions should be used. Administration: Give 6mL If using the 400mg + 200mg vials (or 9 mL if using 3x200mg) intravenously as a slow bolus over 3-5 minutes.	
Water for injections (Supplied with teicoplanin vial)	3.14mL	OD	IV		
Sodium Chloride 0.9%	Flush the cannula with 5mL of sodium chloride 0.9% or the PICC line with 10mL sodium chloride 0.9% before and after each administration via IV route.				

1. SPC Teicoplanin powder for solution for injection/infusion or oral solution. Available from Teicoplanin 400mg powder for solution for injection/infusion or oral solution - Summary of Product Characteristics (SmPC) - (emc) (medicines.org.uk). Last updated 22 Feb 23. 2. Medusa. Teicoplanin. Available from Injectable Medicines Guide - Display - Teicoplanin - Intravenous - Version 14 - IVGuideDisplayMain.asp (wales.nhs.uk). Last updated 04 Oct 2022

Date & time:																				
Given by:																				

Prescriber must be F2 or above, or a suitable non-medical prescriber.									
Signed:		Name: (Print Name)		Professional registration number:		Bleep/ Telephone:		Date:	

