

## Adult Community IV Antibiotic Treatment: Authorisation to Administer and Administration Record – Flucloxacillin 8g B Braun Easypump® II Elastomeric Device

<b>Patient details</b> Name Address  NHS number DOB	<b>Allergies &amp; Intolerances:</b>  No known allergies <input type="checkbox"/> Document nature, details and date of each reaction	<b>Indication for treatment:</b>  Date antibiotic to start in community:  Planned treatment length in community or end date:
--	---	--

<b>eGFR:</b>	<b>Creatinine:</b>	<b>Date:</b>	<b>Weight (kg):</b>	<b>Date:</b>
--------------	--------------------	--------------	---------------------	--------------

Medication	Dose	Frequency	Route	Instructions for preparation and use	Pharmacy check																
Flucloxacillin	8g	Continuous infusion over 24 hours	IV	Reconstitution of vials Using 60mL luer lock syringes, reconstitute each 2g Flucloxacillin vial (x4 vials) with 40mL water for injection. Swirl until dissolved.																	
Water for injections	40mL x4			Order of Additions Prepare the Easypump® as per guidelines.  <b>Total volume= 240ml</b>	<b>Order of additions</b>																
				<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 5%;">1</td> <td style="width: 15%;">Prime</td> <td style="width: 50%;">0.3% Citrate Buffered Sodium Chloride 0.9%</td> <td style="width: 30%;">10mL</td> </tr> <tr> <td>2</td> <td>1<sup>st</sup> Diluent</td> <td>0.3% Citrate Buffered Sodium Chloride 0.9%</td> <td>35mL</td> </tr> <tr> <td>3</td> <td>Drug</td> <td>8g Flucloxacillin</td> <td>160mL</td> </tr> <tr> <td>4</td> <td>2<sup>nd</sup> Diluent</td> <td>0.3% Citrate Buffered Sodium Chloride 0.9%</td> <td>35mL</td> </tr> </table>	1	Prime	0.3% Citrate Buffered Sodium Chloride 0.9%	10mL	2	1 <sup>st</sup> Diluent	0.3% Citrate Buffered Sodium Chloride 0.9%	35mL	3	Drug	8g Flucloxacillin	160mL	4	2 <sup>nd</sup> Diluent	0.3% Citrate Buffered Sodium Chloride 0.9%	35mL	
1	Prime			0.3% Citrate Buffered Sodium Chloride 0.9%	10mL																
2	1 <sup>st</sup> Diluent	0.3% Citrate Buffered Sodium Chloride 0.9%	35mL																		
3	Drug	8g Flucloxacillin	160mL																		
4	2 <sup>nd</sup> Diluent	0.3% Citrate Buffered Sodium Chloride 0.9%	35mL																		
0.3% Citrate Buffered Sodium Chloride 0.9%	As per order of additions	Easypump® = II LT 270-27-S (10mL/h)																			
Sodium Chloride 0.9%	Flush the PICC line with 10mL Sodium Chloride 0.9% before each dose and at the end of the treatment course.																				

1. Medusa Injectable medicines guide (Flucloxacillin) Last updated: 14/7/22 <https://inimed.wales.nhs.uk/IVGuideDisplay.asp> 2. SPC. Flucloxacillin 2g powder for solution for injection vial. Last updated: 30/10/23. <https://www.medicines.org.uk/emc/product/8746/smpc> 3. SOP for Intravenous Infusion Set Flushing – available through NBT LINK/UHBW

Date & time:	/	/	/	/	/	/	/	/	/	/	/	/	/	/	/	/	/	/	/	
Given by:																				

Prescriber must be F2 or above, or a suitable non-medical prescriber.

<b>Signed:</b>		<b>Name: (Print Name)</b>		<b>Professional registration number:</b>		<b>Bleep/ Telephone:</b>		<b>Date:</b>	
----------------	--	---------------------------	--	--	--	--------------------------	--	--------------	--