

## Adult Community IV Antibiotic Treatment: Authorisation to Administer and Administration Record – Flucloxacillin 12g B Braun Easypump® II Elastomeric Device

<b>Patient details</b> Name Address  NHS number DOB		<b>Allergies &amp; Intolerances:</b>  No known allergies <input type="checkbox"/> Document nature, details and date of each reaction		<b>Indication for treatment:</b>  Date antibiotic to start in community:  Planned treatment length in community or end date:						
eGFR:		Creatinine:		Date:		Weight (kg):		Date:		
Medication	Dose	Frequency	Route	Instructions for preparation and use				Pharmacy check		
Flucloxacillin	12g	Continuous infusion over 24 hours	IV	Reconstitution of vials	Using a 20mL luer lock syringe, reconstitute each 1g Flucloxacillin vial (x12vials) with 15mL water for injection. Swirl until dissolved.					
Water for injections	15mL x12			Order of Additions	Prepare the Easypump® as per guidelines. <b>Total volume= 240ml</b>		<b>Order of additions</b>			
				1	Prime	0.3% Citrate Buffered Sodium Chloride 0.9%	10mL			
				2	1 <sup>st</sup> Diluent	0.3% Citrate Buffered Sodium Chloride 0.9%	25mL			
		3	Drug	12g Flucloxacillin	180mL					
		4	2 <sup>nd</sup> Diluent	0.3% Citrate Buffered Sodium Chloride 0.9%	25mL					
0.3% Citrate Buffered Sodium Chloride 0.9%	As per order of additions	Easypump® = II LT 270-27-S (10mL/h)								
Sodium Chloride 0.9%	Flush the PICC line with 10mL Sodium Chloride 0.9% before each dose and at the end of the treatment course.									
1. Medusa Injectable medicines guide (Flucloxacillin) Last updated: 14/7/22 <a href="https://inimed.wales.nhs.uk/IVGuideDisplay.asp">https://inimed.wales.nhs.uk/IVGuideDisplay.asp</a> 2. SPC. Flucloxacillin 1g Powder for Solution for Injection or Infusion Last updated: 10/06/24. <a href="https://www.medicines.org.uk/emc/product/2239/smpc">https://www.medicines.org.uk/emc/product/2239/smpc</a> 3. SOP for Intravenous Infusion Set Flushing – available through NBT LINK/UHBW										
<b>Date &amp; time:</b>										
<b>Given by:</b>										
Prescriber must be F2 or above, or a suitable non-medical prescriber.										
<b>Signed:</b>		<b>Name:</b> (Print Name)		<b>Professional registration number:</b>		<b>Bleep/ Telephone:</b>		<b>Date:</b>		